

Critique of Fermionic $R\nu$ MDM and its Scalar Variants

Krešimir Kumerički, Ivica Picek and Branimir Radovčić

Department of Physics, Faculty of Science, University of Zagreb, P.O.B. 331, HR-10002 Zagreb, Croatia

March 3, 2013

Abstract

We examine the stability of minimal dark matter (MDM) particle-candidates in the setup in which they participate in radiative neutrino ($R\nu$) masses. We first point out the existence of an additional renormalizable term in recently proposed $R\nu$ MDM Lagrangian, which violates the claimed accidental Z_2 symmetry and spoils the stability of the fermionic MDM quintuplet component. We then explore the viability of $R\nu$ MDM variants based on scalar MDM multiplets. There are ubiquitous super-renormalizable terms in the scalar potential which make these scalar multiplets unstable.

PACS: 14.60.Pq; 14.60.St; 95.35.+d; 14.80.Cp

Keywords: Neutrino mass; Exotic leptons; Dark matter; Non-standard scalar fields

1 Introduction

There are two experimental pieces of evidence [1] pressing us to extend the standard model (SM): the evidence for neutrino masses and for the dark matter (DM) content of the Universe. The ongoing experiments at the Large Hadron Collider (LHC) may provide us with new insights on new heavy particles enabling Weinberg's dimension-five operator $LLHH$ [2] as an explanation of the smallness of masses of observed neutrinos.

The known tree level realizations of Weinberg's operator proceed by adding to the particle content of the SM a single, GUT-scale multiplet: a heavy fermion singlet for the type-I [3], a scalar triplet for the type-II [4] and a fermion triplet for the type-III [5] seesaw mechanism. On the other hand additional discrete symmetries are instrumental [6, 7, 8] in bringing the seesaw mechanism to the TeV scale accessible at the LHC.

Alternative lowering of the seesaw scale, without discrete symmetries, can be achieved either by going beyond dimension-five operator or by generating the neutrino masses at the loop level. The approach used by two recent models is to employ non-zero hypercharge seesaw mediators: the dimension-seven mechanism with fermion triplets in [9] and the dimension-nine mechanism with fermion quintuplets in [10, 11]. Both of these mechanisms offer the spectacular triply-charged states verifiable at the LHC.

The approach in [12], which we follow here, is to link a DM candidate to the scale of radiatively generated neutrino ($R\nu$) masses. The authors of [13] exemplify it by employing a 10 TeV scale fermionic quintuplet of zero hypercharge. Let us stress that an isolated $\Sigma \sim (5, 0)$ fermion field or an isolated septuplet $\Phi \sim (7, 0)$ scalar field have been singled out as viable DM particle candidates within the so-called minimal dark matter (MDM) model [14]. The dubbed $R\nu$ MDM model [13] claims to achieve the stability of the neutral MDM quintuplet component without introducing a symmetry beyond the SM gauge symmetries, on account of an accidental Z_2 symmetry arising from the choice of the field content. It has been adopted in more recent paper [15] aimed at successful leptogenesis.

From the point of generating the neutrino masses, the fermion quintuplet has to be accompanied by an appropriate scalar multiplet. High enough scalar field multiplet [13] avoids the terms in the scalar potential which may lead to and induced vacuum expectation value (vev) of the new scalar field, and there are only radiative neutrino masses. This $R\nu$ MDM is realized by accompanying the zero hypercharge quintuplets $\Sigma \sim (5, 0)$ by a hypercharge one sextuplets, $\Phi \sim (6, 1)$, proposing the neutral component Σ^0 as fermionic DM candidate.

In Section 2 we give a brief description of this attempt and explicate a

previously omitted term in the scalar potential, which spoils the stability of Σ^0 particle. In Section 3 we analyze two new seesaw models, the first in which a tentative DM particle appears as a neutral component of the scalar quintuplet $\Phi \sim (5, 0)$, accompanied by a quadruplet seesaw mediator $\Sigma \sim (4, 1)$. In the second model we show that originally viable MDM scalar septuplet $\Phi \sim (7, 0)$ candidate also becomes unstable in the context of generating radiative neutrino masses. We summarize our results in the concluding Section.

2 Instability of fermionic quintuplet $R\nu$ MDM

To allow a Yukawa coupling of the isospin two fermion $\Sigma \sim (5, 0)$ to the lepton doublet $L_L \sim (2, -1)$, necessary for a mass generation mechanism,

$$\mathcal{L}_Y = \overline{L}_L Y \Phi \Sigma + \text{H.c.} , \quad (1)$$

one has to introduce a scalar field Φ of a half-integer ($3/2$ or $5/2$) isospin in addition to the SM higgs $H \sim (2, 1)$. In order to avoid a vev of Φ which might come from quartic $H^3\Phi$ term, the fermionic quintuplet $\Sigma \sim (5, 0)$ has to be accompanied by a sextuplet scalar $\Phi \sim (6, -1)$ as in [13]. These authors have argued that the neutral component of the fermionic quintuplet, Σ^0 , can be a DM candidate on account of an accidental Z_2 symmetry of the Lagrangian. We show that it is a viable DM candidate only if one forbids or finetunes a renormalizable $H\Phi^3$ term with the quartic coupling λ , omitted in [13]. In their tensor notation, the omitted term reads

$$\lambda \Phi^* \Phi^* \Phi H^* + \text{H.c.} , \quad \Phi^* \Phi^* \Phi H^* = \Phi^{*iabcd} \Phi^{*pqrst} \Phi_{abpq} H^{*n} \epsilon_{in} \epsilon_{cs} \epsilon_{dt} , \quad (2)$$

and it can induce a rapid decay of a DM particle at the loop level. For example, the loop diagram on Fig. 1 generates an effective dimension-six operator

$$\mathcal{O}_6 = \overline{L}_L^i \Sigma_{jklm} W_a^l W_b^m H^{*n} \epsilon^{ja} \epsilon^{kb} \epsilon_{in} , \quad (3)$$

which leads to decays of the DM candidate Σ^0 . In particular, we estimate the decay amplitude for $\Sigma^0 \rightarrow \nu W^+ W^- H$ arising from it,

$$A \sim \frac{g^2}{16\pi^2} Y \lambda \frac{m_\Sigma}{m_\Phi^2} \frac{m_\Sigma^2}{m_W^2} . \quad (4)$$

It produces the decay width into four final-state particles,

$$\Gamma \sim \frac{1}{2m_\Sigma} |A|^2 d\text{LIPS}_4 , \quad (5)$$




Figure 1: An example of decays of the heavy lepton Σ at the loop level.

which is given by

$$\Gamma \sim \frac{g^4}{192\pi} \left(\frac{1}{16\pi^2} \right)^4 Y^2 \lambda^2 \frac{m_\Sigma^9}{m_\Phi^4 m_W^4}. \quad (6)$$

The inverse of it, a lifetime of a DM candidate, should be longer than the age of the Universe given by the Hubble time $H^{-1} \approx 10^{17}\text{s}$. For Σ^0 to have the lifetime $\tau_{\text{DM}} > 10^{17}\text{s}$, the quartic coupling λ has to be tiny. By taking $g = 0.65$, $m_\Sigma = m_\Phi = 10 \text{ TeV}$ and $Y = 10^{-1}$, the values adopted from [13], we obtain

$$\lambda < 10^{-20} \quad (7)$$

as an upper limit. In the context of the decaying DM [16] the bound on the τ_{DM} can be nine orders of magnitude larger ($\tau_{\text{DM}} > 10^{26}\text{s}$), implying even stronger limit, $\lambda < 10^{-24}$. This means that Σ^0 is a viable DM candidate only if the coupling in Eq. (2) has an extremely small value. This is in contradiction with the claim in [13], and in order to avoid the fine-tuning one has to impose the Z_2 symmetry forbidding the renormalizable term in Eq. (2), by hand.

Without such discrete symmetry a model more minimal than in [13] is possible with scalar quadruplet $\Phi \sim (4, -1)$ replacing previous sextuplet field. Like in case of previously studied non-zero hypercharge fermion [10, 11], this field Φ develops an induced vev from the quartic ΦH^3 term. The Yukawa terms result in a tree-level dimension-nine seesaw operator. These and additional contributions to the neutrino masses from radiative loop-suppressed diagrams are presented in a separate paper [17].

3 R ν MDM Variants with Scalar MDM

In the light of the described blow to fermionic MDM candidate in the radiative neutrino masses setup, let us now reconsider the scalar MDM multiplets [14, 18, 19]. The models involving hypercharge-zero scalar field in radiatively generating neutrino masses through diagrams displayed on Fig. 2 have two essential features: (i) the scalar field is accompanied with appropriate fermion multiplet; (ii) there are two scalar fields of different hypercharge in play, the one with hypercharge zero being the DM candidate.




Figure 2: Diagram for radiatively generated neutrino masses where one of the scalar fields contains the MDM component.

3.1 Model with scalar quintuplet DM

A minimal possibility to realize the diagram on Fig. 2 through the scalar quintuplet $\Phi_1 \sim (5, 0)$ as MDM candidate is to accompany it with another scalar quintuplet $\Phi_2 \sim (5, 2)$ in conjunction with non-zero hypercharge fermionic quadruplet $\Sigma \sim (4, 1)$ as the seesaw mediator.

With two scalars $\Phi_1 \sim (5, 0)$ and $\Phi_2 \sim (5, 2)$, there are, up to our knowledge previously unconsidered, dimension-three Z_2 noninvariant operators in the Lagrangian

$$\mathcal{L}_S = \mu_1 \Phi_1 \Phi_1 \Phi_1 + \mu_2 \Phi_1 \Phi_2 \Phi_2^* , \quad (8)$$

which in tensor notation read

$$\begin{aligned} \Phi_1 \Phi_1 \Phi_1 &= \Phi_{1ijkl} \Phi_{1mnpq} \Phi_{1rstu} \epsilon^{im} \epsilon^{jn} \epsilon^{kr} \epsilon^{ls} \epsilon^{pt} \epsilon^{qu} , \\ \Phi_1 \Phi_2 \Phi_2^* &= \Phi_{1ijkl} \Phi_{2mnpq} \Phi_2^{*klpq} \epsilon^{im} \epsilon^{jn} . \end{aligned} \quad (9)$$

These terms make the DM candidate Φ_1^0 unstable through loop diagrams like the one on Fig. 3. We estimate the amplitude for the decay $\Phi_1^0 \rightarrow W^+ W^-$

$$A \sim \frac{g^2}{16\pi^2} (\mu_1 + \mu_2) \frac{m_{\Phi_1}^2}{m_W^2} \quad (10)$$

and the decay width

$$\Gamma \sim \frac{g^4}{16\pi} \left(\frac{1}{16\pi^2} \right)^2 (\mu_1 + \mu_2)^2 \frac{m_{\Phi_1}^3}{m_W^4} . \quad (11)$$

To have the lifetime of a DM candidate $\tau_{\text{DM}} > 10^{17}\text{s}$, the parameters $\mu_{1,2}$ in Eq. (8) have to have very small values, $\mu_{1,2} < 10^{-10}\text{eV}$.




Figure 3: An example of the loop diagram for quintuplet Φ_1 decays.

3.2 Model with scalar septuplet DM

The second possibility to realize the diagram on Fig. 2 is through the scalar septuplet $\Phi_1 \sim (7, 0)$ as MDM, together with another scalar septuplet $\Phi_2 \sim (7, 2)$, now in conjunction with fermionic sextuplet $\Sigma \sim (6, 1)$.

For a single scalar septuplet Φ_1 , the term Φ_1^3 is forbidden by Bose statistics [18], but in presence of another septuplet Φ_2 there is the dimension-three Z_2 noninvariant operator

$$\begin{aligned} \mathcal{L}_S &= \mu \Phi_1 \Phi_2 \Phi_2^* , \\ \Phi_1 \Phi_2 \Phi_2^* &= \Phi_{1ijklmn} \Phi_{2pqrstu} \Phi_2^{lmnstu} \epsilon^{ip} \epsilon^{jq} \epsilon^{kr} , \end{aligned} \quad (12)$$

making the DM candidate Φ_1^0 unstable. These terms make the DM candidate Φ_1^0 unstable through loop diagrams, like the one on Fig. 4. We estimate the amplitude for the decay $\Phi_1^0 \rightarrow W^+ W^- Z^0$

$$A \sim \frac{g^3}{16\pi^2} \frac{\mu}{m_{\Phi_1}} \frac{m_{\Phi_1}^3}{m_W^3} , \quad (13)$$

and the decay width

$$\Gamma \sim \frac{g^6}{32\pi} \left(\frac{1}{16\pi^2} \right)^3 \mu^2 \frac{m_{\Phi_1}^5}{m_W^6} . \quad (14)$$




Figure 4: An example of the loop diagram for septuplet Φ_1 decays.

To have the lifetime of a DM candidate $\tau_{\text{DM}} > 10^{17}\text{s}$, the parameter μ in Eq. (12) is restricted to very small value, $\mu < 10^{-11}\text{eV}$.

To conclude, a major blow to scalar R ν MDM variants comes from super-renormalizable dimension-three Φ^3 operators. Besides these renormalizable Z_2 violating operators, there are possible effects from the Planck scale considered in [14, 16] and references therein.

4 Conclusions

We address in a common framework the open questions of the neutrino masses and the dark matter content of the Universe. In view of the immense DM possibilities, a minimality of the DM model appears as a criterion essential for its predictivity and testability. The minimality is satisfied by adding only one extra $SU(2)_L$ multiplet in the MDM setup. However, by imposing additional seesaw mission to a given MDM multiplet, it has to be accompanied by additional scalar field introducing new obstacles to the DM stability. It would be appealing if the DM candidate, a neutral component of a certain higher weak-isospin multiplet, could be stabilized without introducing a symmetry beyond the SM gauge symmetries, as claimed in [13]. We have shown that this is not possible for already selected MDM particles, if the neutrino masses are generated through simple one-loop diagrams on Fig. 2. The obstacle for the fermionic quintuplet MDM candidate in [13] comes from renormalizable quartic term omitted there. Concerning the scalar R ν MDM variants, we have shown that the stability of the scalar quintuplet and septuplet MDM candidate is violated by ubiquitous super-renormalizable terms that can be tamed only by imposing extra discrete symmetry.

References

- [1] K. Nakamura *et al.* (Particle Data Group), *Review of particle physics*, J. Phys. G **37**, 075021 (2010) [<http://pdg.lbl.gov/>].
- [2] S. Weinberg, *Baryon- and Lepton-Nonconserving Processes*, Phys. Rev. Lett. **43**, 1566 (1979).
- [3] P. Minkowski, *$\mu \rightarrow e \gamma$ At A Rate Of One Out Of 1-Billion Muon Decays?*, Phys. Lett. B **67**, 421 (1977); T. Yanagida, in *Proceedings of the Workshop on Unified Theory and the Baryon Number of the Universe*, eds. O. Sawada and A. Sugamoto (KEK, Tsukuba, 1979), p. 95; M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Freedman (North Holland, Amsterdam, 1979), p. 315; S.L. Glashow, in *Quarks and Leptons*, edited by M. Lévy *et al.* (Plenum, New York, 1980), p. 707; R.N. Mohapatra and G. Senjanović, *Neutrino Mass and Spontaneous Parity Nonconservation*, Phys. Rev. Lett. **44**, 912 (1980).
- [4] W. Konetschny and W. Kummer, *Nonconservation of total lepton number with scalar bosons*, Phys. Lett. B **70**, 433 (1977); M. Magg and C. Wetterich, *Neutrino mass problem and gauge hierarchy*, Phys. Lett. B **94**, 61 (1980); J. Schechter and J.W.F. Valle, *Neutrino masses in $SU(2) \times U(1)$ theories*, Phys. Rev. D **22**, 2227 (1980); T.P. Cheng and L.F. Li, *Neutrino masses, mixings, and oscillations in $SU(2) \times U(1)$ models of electroweak interactions*, Phys. Rev. D **22**, 2860 (1980); G. Lazarides, Q. Shafi, and C. Wetterich, *Proton lifetime and fermion masses in an $SO(10)$ model*, Nucl. Phys. B **181**, 287 (1981); R.N. Mohapatra and G. Senjanović, *Neutrino masses and mixings in gauge models with spontaneous parity violation*, Phys. Rev. D **23**, 165 (1981).
- [5] R. Foot, H. Lew, X.G. He and G.C. Joshi, *Seesaw neutrino masses induced by a triplet of leptons*, Z. Phys. C **44**, 441 (1989).
- [6] E. Ma, *Naturally small seesaw neutrino mass with no new physics beyond the TeV scale*, Phys. Rev. Lett. **86**, 2502 (2001) [[hep-ph/0011121](#)].
- [7] W. Grimus, L. Lavoura and B. Radovčić, *Type II seesaw mechanism for Higgs doublets and the scale of new physics*, Phys. Lett. B **674**, 117 (2009) [[0902.2325](#) [[hep-ph](#)]].
- [8] Z.Z. Xing and S. Zhou *Multiple seesaw mechanisms of neutrino masses at the TeV scale*, Phys. Lett. B **679**, 249 (2009) [[0906.1757](#) [[hep-ph](#)]].

- [9] K.S. Babu, S. Nandi and Z. Tavartkiladze, *New Mechanism for Neutrino Mass Generation and Triply Charged Higgs Bosons at the LHC*, Phys. Rev. D **80**, 071702 (2009) [0905.2710 [hep-ph]].
- [10] I. Picek and B. Radovčić, *Novel TeV-scale seesaw mechanism with Dirac mediators*, Phys. Lett. B **687**, 338 (2010) [0911.1374 [hep-ph]].
- [11] K. Kumerički, I. Picek and B. Radovčić, *Exotic Seesaw-Motivated Heavy Leptons at the LHC*, Phys. Rev. D **84**, 093002 (2011) [1106.1069 [hep-ph]].
- [12] E. Ma, *Verifiable radiative seesaw mechanism of neutrino mass and dark matter*, Phys. Rev. D **73**, 077301 (2006) [hep-ph/0601225].
- [13] Y. Cai, X-G He, M. Ramsey-Musolf and L-H Tsai, *$R\nu$ MDM and Lepton Flavor Violation*, JHEP 1112, 054 (2011) (2011)[1108.0969 [hep-ph]].
- [14] M. Cirelli, N. Fornengo and A. Strumia, *Minimal dark matter*, Nucl. Phys. B **753**, 178 (2006) [hep-ph/0512090]; M. Cirelli and A. Strumia, *Minimal Dark Matter: model and results*, Phys. Rev. D **80**, 071702 (2009) [0905.2710 [hep-ph]].
- [15] C-H Chen and S.S.C. Law, *Exotic fermion multiplets as a solution to baryon asymmetry, dark matter and neutrino masses*, Phys. Rev. D **85**, 055012 (2012) [1111.5462 [hep-ph]].
- [16] M. S. Boucenna, R. A. Lineros and J. W. F. Valle *Planck-scale effects on WIMP dark matter*, [1204.2576 [hep-ph]].
- [17] K. Kumerički, I. Picek and B. Radovčić, *TeV-scale Seesaw with Quintuplet Fermions*, Phys. Rev. D **86**, 013006 (2012) [1204.6599 [hep-ph]].
- [18] T. Hambye, F.-S. Ling, L. Lopez Honorez and J. *Scalar Multiplet Dark Matter*, JHEP 0907, 090 (2009), Erratum-ibid. 1005, 066 (2010) [0903.4010 [hep-ph]]
- [19] K. Hally, H.E. Logan and T. Pilkington, *Constraints on large scalar multiplets from perturbative unitarity*, Phys. Rev. D **85**, 095017 (2012) [1202.5073 [hep-ph]].